SAN BERNARDINO UNIFIED SCHOOL DISTRICT, CA

MIDDLE SCHOOL YOUTH RISK BEHAVIOR SURVEY
2003

Table 2I. Percentage of Middle School Students Who Had Rarely or Never Worn Seatbelts* and Who Had Ever Ridden in a Car with a Driver Who Had Been Drinking Alcohol, by Sex, Race/ Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | 1ale | Female | | Total | |
|-----------------------------------|-------------|---------------------|--------|--------|-------|--------|
| Question | % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Rarely or never wore seat belts | | | | | | |
| Race/Ethnicity | | | | | | |
| White [§] | 10.1 | 8.7 | 6.2 | 5.0 | 8.1 | 4.9 |
| Black§ | 4.0 | 3.3 | 6.6 | 3.9 | 5.4 | 2.5 |
| Hispanic | 13.3 | 3.3 | 10.4 | 3.0 | 11.9 | 2.6 |
| Other | 9.6 | 5.9 | 4.3 | 3.5 | 7.3 | 3.3 |
| Grade | | | | | | |
| 6 | 8.1 | 4.4 | 2.8 | 3.7 | 5.4 | 2.8 |
| 7 | 10.8 | 3.5 | 7.7 | 2.6 | 9.3 | 2.4 |
| 8 | 13.1 | 3.9 | 14.8 | 4.5 | 13.9 | 3.3 |
| Total | 10.8 | 2.3 | 8.5 | 2.4 | 9.6 | 1.8 |
| Rode with a driver who had been o | rinking alc | ohol | | | | |
| Race/Ethnicity | | | | | | |
| White | 21.9 | 7.7 | 25.4 | 11.0 | 23.5 | 6.9 |
| Black | 28.4 | 8.1 | 29.8 | 8.8 | 29.1 | 6.9 |
| Hispanic | 33.5 | 5.3 | 33.6 | 4.7 | 33.6 | 4.1 |
| Other | 26.9 | 8.0 | 31.5 | 8.0 | 29.8 | 5.9 |
| Grade | | | | | | |
| 6 | 18.9 | 7.0 | 22.4 | 5.1 | 20.8 | 5.1 |
| 7 | 28.7 | 5.4 | 28.3 | 6.0 | 28.5 | 4.4 |
| 8 | 42.5 | 7.2 | 45.8 | 7.1 | 44.1 | 5.5 |
| Total | 29.9 | 4.8 | 31.4 | 4.4 | 30.7 | 3.9 |

^{*}When riding in a car.

^{†95%} confidence interval.

[§]Non-Hispanic.

Table 3I. Percentage of Middle School Students Who Had Rarely or Never Worn Bicycle Helmets* or Rollerblade or Skateboard Helmets,† by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Fe | Female | | otal |
|---------------------------|----------------------|-------------------|---------------|--------|------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Rarely or never wore bicy | cle helmets | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 66.3 | 9.6 | 50.7 | 12.9 | 59.7 | 7.8 |
| Black ¹ | 70.4 | 11.1 | 75.0 | 9.3 | 72.6 | 8.2 |
| Hispanic | 75.4 | 3.8 | 65.2 | 6.4 | 71.2 | 3.4 |
| Other | 66.9 | 10.8 | ** | _ | 64.2 | 9.8 |
| Grade | | | | | | |
| 6 | 71.3 | 8.9 | 55.8 | 8.5 | 64.5 | 6.8 |
| 7 | 66.9 | 5.7 | 59.1 | 7.4 | 63.5 | 5.2 |
| 8 | 78.9 | 6.3 | 80.4 | 9.0 | 79.5 | 5.1 |
| Total | 72.1 | 4.3 | 64.5 | 5.5 | 68.8 | 3.7 |
| Rarely or never wore a he | elmet when using rol | lerblades or ridi | ng a skateboa | rd | | |
| Race/Ethnicity | | | | | | |
| White | 73.1 | 9.7 | 67.6 | 10.5 | 70.6 | 7.2 |
| Black | _ | _ | _ | _ | 70.3 | 9.7 |
| Hispanic | 75.5 | 5.4 | 73.5 | 5.9 | 74.7 | 4.5 |
| Other | _ | _ | _ | _ | 70.8 | 9.2 |
| Grade | | | | | | |
| 6 | 70.0 | 10.1 | 69.5 | 8.9 | 70.0 | 8.8 |
| 7 | 69.7 | 6.0 | 69.0 | 7.4 | 69.3 | 4.6 |
| 8 | 81.0 | 6.7 | 79.6 | 9.2 | 80.4 | 6.3 |
| Total | 73.3 | 4.6 | 72. I | 4.8 | 72.8 | 3.9 |

^{*}Among the 75.2% of students who had ridden a bicycle.

[†]Among the 60.1% of students who had used rollerblades or ridden a skateboard.

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 4I. Percentage of Middle School Students Who Had Ever Carried a Weapon,* Who Had Ever Been in a Physical Fight, and Who Had Ever Been Hurt in a Physical Fight,† by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | 1ale | Female | | Total | |
|-------------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Carried a weapon | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 43.6 | 11.3 | 10.9 | 7.0 | 27.5 | 7.0 |
| Black [¶] | 36.4 | 9.0 | 12.1 | 5.3 | 23.9 | 5.7 |
| Hispanic | 35.9 | 5.1 | 17.5 | 4.9 | 27.0 | 4.1 |
| Other | 31.5 | 8.8 | 14.5 | 7.4 | 22.9 | 6.1 |
| Grade | | | | | | |
| 6 | 25.9 | 6.1 | 9.1 | 7.0 | 17.5 | 6.0 |
| 7 | 37.4 | 6.5 | 13.7 | 3.5 | 25.8 | 4.8 |
| 8 | 49.6 | 7.0 | 21.1 | 5.8 | 35.7 | 5.5 |
| Total | 37.4 | 4.6 | 14.8 | 3.5 | 26.2 | 3.6 |
| Were in a physical fight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 62.7 | 12.3 | 36.3 | 10.9 | 50.0 | 8.8 |
| Black | 80.6 | 7.3 | 54.3 | 11.0 | 67.2 | 6.8 |
| Hispanic | 69.1 | 5.3 | 42.3 | 5.5 | 56.1 | 3.9 |
| Other | 70.1 | 8.4 | 39.9 | 10.8 | 55.0 | 6.4 |
| Grade | | | | | | |
| 6 | 67.9 | 7.2 | 34.1 | 9.0 | 51.4 | 4.9 |
| 7 | 68.1 | 5.7 | 43.2 | 6.9 | 55.9 | 5.2 |
| 8 | 74.3 | 5.9 | 53.5 | 6.1 | 64.1 | 4.8 |
| Total | 69.8 | 3.8 | 43.6 | 4.5 | 57.0 | 3.1 |
| Were hurt in a physical fight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 7.4 | 4.2 | 3.6 | 3.5 | 5.5 | 2.8 |
| Black | 5.8 | 3.6 | 3.5 | 3.0 | 4.6 | 2.3 |
| Hispanic | 7.9 | 2.8 | 3.2 | 1.4 | 5.6 | 1.5 |
| Other | 10.8 | 6.9 | 2.7 | 2.8 | 6.8 | 4.0 |
| Grade | | | | | | |
| 6 | 6.5 | 3.5 | 2.5 | 2.0 | 4.5 | 2.1 |
| 7 | 8.8 | 2.8 | 2.2 | 1.8 | 5.6 | 2.0 |
| 8 | 6.4 | 2.8 | 5.4 | 2.6 | 5.9 | 1.8 |
| Total | 7.4 | 1.8 | 3.3 | 1.3 | 5.4 | 1.2 |

^{*}For example, a gun, knife, or club.

[†]Had to be treated by a doctor or nurse.

^{§95%} confidence interval.

¹Non-Hispanic.

Table 5I. Percentage of Middle School Students Who Had Ever Seriously Thought About Killing Themselves, Who Had Ever Made a Plan About How To Kill Themselves, and Who Had Ever Tried To Kill Themselves, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Fe | male | Total | |
|---------------------------------|------------|---------|------|--------------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Seriously thought about killing | themselves | | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 17.1 | 6.7 | 25.3 | 9.6 | 20.9 | 6.3 |
| Black [†] | 25.5 | 9.2 | 24.1 | 7.9 | 24.8 | 6.3 |
| Hispanic | 18.0 | 3.7 | 28.3 | 5.2 | 23.0 | 3.3 |
| Other | 25.9 | 8.1 | 26.0 | 8.5 | 25.6 | 5.9 |
| Grade | | | | | | |
| 6 | 16.6 | 6.6 | 19.8 | 6.8 | 18.2 | 4.0 |
| 7 | 20.3 | 3.5 | 23.5 | 5.0 | 21.9 | 3.1 |
| 8 | 21.2 | 5.6 | 37.0 | 7.7 | 28.8 | 4.6 |
| Total | 19.5 | 3.2 | 26.7 | 4 . l | 23.1 | 2.5 |
| Made a plan about how to kill t | hemselves | | | | | |
| Race/Ethnicity | | | | | | |
| White | 11.6 | 6.0 | 17.2 | 7.6 | 14.3 | 5.0 |
| Black | 9.7 | 5.8 | 8.6 | 4.5 | 9.1 | 3.3 |
| Hispanic | 10.0 | 2.6 | 13.2 | 3.6 | 11.6 | 2.0 |
| Other | 11.1 | 5.6 | 13.4 | 7.4 | 12.0 | 4.4 |
| Grade | | | | | | |
| 6 | 8.7 | 5.8 | 7.0 | 5.5 | 8.0 | 3.0 |
| 7 | 10.2 | 3.0 | 10.2 | 2.9 | 10.2 | 2.3 |
| 8 | 11.6 | 3.8 | 19.9 | 5.4 | 15.7 | 2.9 |
| Total | 10.2 | 2.5 | 12.7 | 3.1 | 11.5 | 1.9 |
| Tried to kill themselves | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 6.6 | 5.0 | 14.8 | 8.3 | 10.6 | 4.6 |
| Black | 15.5 | 8.8 | 8.3 | 4.4 | 11.8 | 4.7 |
| Hispanic | 7.7 | 3.0 | 10.4 | 3.8 | 9.0 | 2.3 |
| Other | 12.1 | 7.5 | 11.4 | 5.3 | 11.9 | 4.2 |
| Grade | | | | | | |
| 6 | 7.9 | 6.1 | 1.4 | 1.4 | 4.7 | 3.6 |
| 7 | 10.4 | 3.5 | 9.4 | 2.8 | 9.9 | 2.5 |
| 8 | 8.3 | 3.7 | 20.7 | 6.9 | 14.3 | 4.4 |
| Total | 9.2 | 2.7 | 10.7 | 3.3 | 9.9 | 2.3 |

^{*95%} confidence interval.

[†]Non-Hispanic.

Table 6I. Percentage of Middle School Students Who Had Ever Smoked Cigarettes, by Sex, Race/ Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|---|------|---------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Lifetime cigarette use [†] | | | | | | |
| Race/Ethnicity | | | | | | |
| White§ | 28.6 | 9.3 | 26.6 | 8.7 | 28.1 | 7.2 |
| Black [§] | 33.2 | 9.7 | 42.3 | 12.5 | 37.9 | 7.4 |
| Hispanic | 32.3 | 6.2 | 29.4 | 6.5 | 31.0 | 5.2 |
| Other | 25.7 | 9.0 | 31.8 | 9.1 | 28.2 | 7.8 |
| Grade | | | | | | |
| 6 | 18.6 | 8.0 | 17.6 | 6.3 | 18.5 | 3.0 |
| 7 | 29.6 | 6.6 | 30.5 | 7.0 | 30.0 | 5.4 |
| 8 | 48.2 | 8.6 | 46.2 | 8.2 | 47.2 | 7.0 |
| Total | 31.5 | 5.5 | 31.4 | 5.2 | 31.5 | 4.3 |
| Lifetime daily cigarette use ¹ | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 3.0 | 2.3 | 3.1 | 2.4 | 3.0 | 1.5 |
| Black | 4.9 | 4.2 | 5.0 | 4.7 | 4.9 | 2.9 |
| Hispanic | 3.7 | 2.2 | 2.5 | 1.3 | 3.1 | 1.4 |
| Other | 5.9 | 3.8 | 1.2 | 1.8 | 3.7 | 2.3 |
| Grade | | | | | | |
| 6 | 2.4 | 2.1 | 1.9 | 2.4 | 2.1 | 1.4 |
| 7 | 4.1 | 2.2 | 4.7 | 2.4 | 4.4 | 1.8 |
| 8 | 4.7 | 3.5 | 2.7 | 1.8 | 3.8 | 2.3 |
| Total | 3.8 | 1.7 | 3.1 | 1.3 | 3.5 | 1.1 |

^{*95%} confidence interval.

[†]Ever tried cigarette smoking, even one or two puffs.

[§]Non-Hispanic.

¹Ever smoked at least one cigarette every day for 30 days.

Table 7I. Percentage of Middle School Students Who Smoked Cigarettes, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | 1 ale | Female | | Total | |
|---|------|--------------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Current cigarette use [†] | | | | | | |
| Race/Ethnicity | | | | | | |
| White§ | 2.9 | 2.6 | 11.0 | 6.7 | 6.8 | 3.7 |
| Black [§] | 7.4 | 4.8 | 6.3 | 4.7 | 6.8 | 3.5 |
| Hispanic | 8.1 | 4.1 | 8.4 | 3.1 | 8.2 | 2.9 |
| Other | 5.0 | 3.7 | 4.3 | 3.6 | 4.6 | 2.6 |
| Grade | | | | | | |
| 6 | 3.1 | 3.6 | 4.2 | 2.6 | 3.6 | 2.3 |
| 7 | 6.7 | 2.7 | 8.5 | 4.9 | 7.6 | 3.1 |
| 8 | 11.0 | 6.0 | 10.7 | 6.5 | 10.9 | 5.7 |
| Total | 6.9 | 2.8 | 8.3 | 3.0 | 7.5 | 2.5 |
| Current frequent cigarette use ¹ | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 0.9 | 1.2 | 0.9 | 1.2 | 0.9 | 0.9 |
| Black | 1.1 | 2.1 | 0.0 | 0.0 | 0.5 | 1.1 |
| Hispanic | 2.0 | 2.9 | 0.6 | 0.7 | 1.3 | 1.6 |
| Other | 1.8 | 2.4 | 0.0 | 0.0 | 0.9 | 1.3 |
| Grade | | | | | | |
| 6 | 2.0 | 3.6 | 0.2 | 0.3 | 1.1 | 2.0 |
| 7 | 0.2 | 0.4 | 0.4 | 0.5 | 0.3 | 0.3 |
| 8 | 2.1 | 2.5 | 1.0 | 1.2 | 1.6 | 1.8 |
| Total | 1.6 | 1.8 | 0.5 | 0.4 | 1.1 | 1.1 |

^{*95%} confidence interval.

 $^{^{\}dagger}\text{Smoked}$ cigarettes on ≥ 1 of the 30 days preceding the survey.

[§]Non-Hispanic.

 $^{^{1}}$ Smoked cigarettes on \geq 20 of the 30 days preceding the survey.

Table 8I. Percentage of Middle School Students Who Used Smokeless Tobacco,* Smoked Cigars,† and Used Any Tobacco Product,§ by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | N | 1ale | Female | | Total | |
|-------------------------------|------|---------------------|--------|--------|-------|--------|
| Question | % | CI ¹ (±) | % | CI (±) | % | CI (±) |
| Current smokeless tobacco use | | | | | | |
| Race/Ethnicity | | | | | | |
| White** | 1.2 | 1.6 | 4.5 | 4.3 | 2.8 | 2.3 |
| Black** | 3.4 | 3.5 | 1.2 | 2.3 | 2.3 | 2.1 |
| Hispanic | 4.2 | 2.2 | 4.3 | 1.6 | 4.2 | 1.5 |
| Other | 2.9 | 3.0 | 3.9 | 3.6 | 3.3 | 2.9 |
| Grade | | | | | | |
| 6 | 1.6 | 1.9 | 2.4 | 2.3 | 2.0 | 1.4 |
| 7 | 3.7 | 2.1 | 3.9 | 2.1 | 3.8 | 1.5 |
| 8 | 4.8 | 3.9 | 3.9 | 2.0 | 4.4 | 2.3 |
| Total | 3.4 | 1.7 | 3.6 | 1.3 | 3.5 | 1.2 |
| Current cigar use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 2.8 | 2.6 | 4.4 | 3.2 | 3.6 | 1.9 |
| Black | 4.5 | 3.9 | 7.2 | 5.0 | 5.9 | 3.0 |
| Hispanic | 7.8 | 2.8 | 6.1 | 2.1 | 7.1 | 1.9 |
| Other | 4.9 | 3.6 | 4.4 | 3.8 | 5.0 | 2.7 |
| Grade | | | | | | |
| 6 | 2.9 | 2.7 | 2.8 | 2.3 | 3.1 | 1.5 |
| 7 | 6.5 | 2.9 | 7.4 | 3.6 | 7.0 | 2.3 |
| 8 | 9.2 | 3.6 | 7.5 | 2.7 | 8.3 | 2.3 |
| Total | 6.2 | 1.9 | 6.0 | 1.8 | 6.2 | 1.3 |
| Current tobacco use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 4.4 | 3.5 | 13.9 | 7.6 | 8.9 | 4.1 |
| Black | 9.2 | 5.1 | 7.3 | 4.9 | 8.3 | 3.7 |
| Hispanic | 9.3 | 3.8 | 9.3 | 3.1 | 9.3 | 2.9 |
| Other | 6.2 | 4.0 | 6.5 | 4.7 | 6.3 | 3.4 |
| Grade | | | | | | |
| 6 | 2.9 | 2.4 | 5.0 | 3.0 | 3.9 | 1.9 |
| 7 | 8.4 | 3.0 | 10.4 | 5.0 | 9.4 | 3.2 |
| 8 | 13.8 | 6.3 | 12.0 | 5.9 | 13.0 | 5.4 |
| Total | 8.3 | 2.8 | 9.5 | 3.0 | 8.9 | 2.5 |

^{*}Used chewing tobacco, snuff, or dip on ≥ 1 of the 30 days preceding the survey.

 $^{^{\}dagger}$ Smoked cigars, cigarillos, or little cigars on ≥ 1 of the 30 days preceding the survey.

[§]Reported current cigarette use, current cigar use, or had used chewing tobacco, snuff, or dip on ≥ 1 of the 30 days preceding the survey.

^{195%} confidence interval.

^{**}Non-Hispanic.

Table 9I. Percentage of Middle School Students Who Had Ever Had a Drink of Alcohol,* Who Had Ever Used Marijuana, and Who Had Ever Used Cocaine,† by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Lifetime alcohol use | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 36.6 | 10.4 | 32.7 | 10.8 | 34.7 | 8.0 |
| Black [¶] | 38.2 | 11.1 | 33.4 | 9.5 | 35.7 | 6.6 |
| Hispanic | 46.4 | 6.6 | 44.7 | 6.2 | 45.6 | 5.0 |
| Other | 35.4 | 9.5 | 38.1 | 10.3 | 37.2 | 7.3 |
| Grade | | | | | | |
| 6 | 28.1 | 9.3 | 27.0 | 9.9 | 27.7 | 7.4 |
| 7 | 43.1 | 5.3 | 36.5 | 7.1 | 39.9 | 5.1 |
| 8 | 55.9 | 6.3 | 55.7 | 8.5 | 55.8 | 6.1 |
| Total | 42.5 | 4.5 | 39.6 | 5.4 | 41.1 | 4.1 |
| Lifetime marijuana use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 10.8 | 6.6 | 9.7 | 5.9 | 10.2 | 4.4 |
| Black | 23.3 | 7.6 | 12.5 | 6.6 | 17.8 | 5.2 |
| Hispanic | 18.6 | 4.3 | 13.5 | 4.3 | 16.1 | 3.5 |
| Other | 16.4 | 7.8 | 10.5 | 6.7 | 13.3 | 6.0 |
| Grade | | | | | | |
| 6 | 13.8 | 7.7 | 1.4 | 1.3 | 7.6 | 4.3 |
| 7 | 15.3 | 3.7 | 12.3 | 4.0 | 13.9 | 2.8 |
| 8 | 25.0 | 6.9 | 21.9 | 6.3 | 23.5 | 5.0 |
| Total | 17.9 | 4.0 | 12.4 | 3.4 | 15.2 | 2.9 |
| Lifetime cocaine use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 2.5 | 2.5 | 2.1 | 2.2 | 2.3 | 1.6 |
| Black | 1.7 | 2.4 | 1.1 | 1.6 | 1.4 | 1.4 |
| Hispanic | 4.4 | 2.0 | 6.0 | 2.5 | 5.2 | 1.7 |
| Other | 3.3 | 2.5 | 2.2 | 2.0 | 2.7 | 1.7 |
| Grade | | | | | | |
| 6 | 1.7 | 1.9 | 1.1 | 1.3 | 1.4 | 1.2 |
| 7 | 3.9 | 2.4 | 4.1 | 1.8 | 4.0 | 1.6 |
| 8 | 4.4 | 2.3 | 7.3 | 3.6 | 5.8 | 2.3 |
| Total | 3.5 | 1.3 | 4.1 | 1.6 | 3.8 | 1.1 |

^{*}Ever had a drink of alcohol, other than a few sips.

 $^{^{\}dagger}\textsc{Ever}$ used any form of cocaine (e.g., powder, "crack," or "freebase").

^{§95%} confidence interval.

¹Non-Hispanic.

Table 10l. Percentage of Middle School Students Who Had Ever Used Inhalants* and Who Had Ever Used Steroids, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|-----------------------|------|---------------------|--------|--------|-------|--------|
| Question | % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Lifetime inhalant use | | | | | | |
| Race/Ethnicity | | | | | | |
| White [§] | 10.0 | 5.1 | 11.6 | 6.1 | 10.7 | 4.4 |
| Black [§] | 7.9 | 5.0 | 5.1 | 3.4 | 6.5 | 2.9 |
| Hispanic | 12.2 | 3.3 | 15.7 | 4.1 | 13.9 | 2.8 |
| Other | 8.2 | 5.1 | 11.3 | 5.6 | 9.6 | 3.4 |
| Grade | | | | | | |
| 6 | 6.7 | 4.0 | 7.9 | 6. l | 7.2 | 3.4 |
| 7 | 12.0 | 3.7 | 9.1 | 3.1 | 10.6 | 2.5 |
| 8 | 13.6 | 4.0 | 19.6 | 5.6 | 16.5 | 3.4 |
| Total | 10.7 | 2.4 | 12.6 | 3.1 | 11.6 | 2.2 |
| ifetime steroid use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 4.3 | 3.2 | 4.9 | 4.9 | 4.6 | 2.9 |
| Black | 5.2 | 4.1 | 1.2 | 1.7 | 3.2 | 2.2 |
| Hispanic | 4.2 | 2.2 | 2.8 | 2.1 | 3.5 | 2.0 |
| Other | 3.4 | 3.0 | 2.8 | 3.2 | 3.1 | 2.1 |
| Grade | | | | | | |
| 6 | 2.2 | 2.0 | 1.3 | 2.1 | 1.7 | 1.9 |
| 7 | 6.1 | 2.6 | 3.0 | 1.9 | 4.6 | 1.6 |
| 8 | 4.4 | 2.4 | 3.1 | 2.6 | 3.7 | 2.2 |
| Total | 4.3 | 1.5 | 2.8 | 1.5 | 3.6 | 1.3 |

^{*}Sniffed glue, breathed the contents of spray cans, or inhaled any paints or sprays to get high.

^{†95%} confidence interval.

[§]Non-Hispanic.

Table III. Percentage of Middle School Students Who Had Initiated Drug-Related Behaviors Before Age II Years, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | | Male | | Female | | Total | |
|----------------------------|-----------------------|---------|------|--------|------|--------|--|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) | |
| Smoked a whole cigarett | e before age 11 years | S | | | | | |
| Race/Ethnicity | | | | | | | |
| White [†] | 3.4 | 2.3 | 3.6 | 3.3 | 3.5 | 1.9 | |
| Black [†] | 7.8 | 5.7 | 5.6 | 4.0 | 6.7 | 3.4 | |
| Hispanic | 8.3 | 3.5 | 5.5 | 2.2 | 7.1 | 2.1 | |
| Other | 8.6 | 5.2 | 5.7 | 3.7 | 7.2 | 2.9 | |
| Grade | | | | | | | |
| 6 | 5.5 | 6.1 | 2.8 | 2.2 | 4.4 | 3.0 | |
| 7 | 8.7 | 3.9 | 5.2 | 2.4 | 7.0 | 2.7 | |
| 8 | 7.3 | 2.6 | 7.9 | 3.8 | 7.6 | 2.3 | |
| Total | 7.2 | 2.7 | 5.2 | 1.6 | 6.3 | 1.6 | |
| Had first drink of alcohol | before age 11 years§ | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 17.5 | 7.0 | 19.1 | 9.2 | 18.3 | 6.5 | |
| Black | 20.3 | 8.0 | 14.3 | 8.6 | 17.3 | 5.8 | |
| Hispanic | 25.7 | 4.2 | 21.1 | 4.5 | 23.5 | 3.4 | |
| Other | 17.9 | 6.9 | 20.8 | 8.4 | 19.3 | 5.4 | |
| Grade | | | | | | | |
| 6 | 16.2 | 5.3 | 18.5 | 7.0 | 17.5 | 4.6 | |
| 7 | 24.7 | 4.4 | 18.0 | 5.6 | 21.5 | 4.3 | |
| 8 | 26.7 | 4.6 | 19.3 | 6.1 | 23.1 | 4.6 | |
| Total | 22.7 | 2.8 | 19.0 | 3.6 | 20.9 | 2.5 | |
| Tried marijuana before ag | ge II years | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 2.4 | 3.0 | 0.0 | 0.0 | 1.2 | 1.5 | |
| Black | 7.9 | 5.2 | 2.9 | 2.5 | 5.3 | 3.0 | |
| Hispanic | 4.8 | 2.0 | 1.3 | 0.8 | 3.1 | 1.2 | |
| Other | 3.9 | 3.8 | 1.8 | 2.5 | 2.8 | 2.3 | |
| Grade | | | | | | | |
| 6 | 3.4 | 3.2 | 0.3 | 0.4 | 1.9 | 1.7 | |
| 7 | 6.1 | 2.7 | 2.8 | 2.0 | 4.5 | 2.1 | |
| 8 | 4.8 | 2.2 | 1.1 | 1.1 | 3.0 | 1.4 | |
| Total | 4.9 | 1.7 | 1.4 | 0.8 | 3.2 | 1.1 | |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Other than a few sips.

Table 12I. Percentage of Middle School Students Who Had Engaged in Sexual Behaviors, by Sex, Race/ Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|-------------------------------------|--------------|---------|--------|--------------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Had ever had sexual intercourse | | | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 17.6 | 8.9 | 8.6 | 6.9 | 12.9 | 5.7 |
| Black [†] | 26.1 | 9.2 | 13.7 | 7.6 | 19.6 | 5.9 |
| Hispanic | 21.5 | 4.4 | 8.4 | 2.7 | 15.0 | 2.8 |
| Other | 14.0 | 6.8 | 9.9 | 5.1 | 11.8 | 4.5 |
| Grade | | | | | | |
| 6 | 14.5 | 7.6 | 5.8 | 3.7 | 10.0 | 4.4 |
| 7 | 17.3 | 4.7 | 7.7 | 3.1 | 12.6 | 3.4 |
| 8 | 32.9 | 5.8 | 14.6 | 4.2 | 23.7 | 3.5 |
| Total | 21.3 | 4.2 | 9.5 | 2.7 | 15.4 | 2.7 |
| lad first sexual intercourse before | age II yea | ırs | | | | |
| Race/Ethnicity | | | | | | |
| White | 2.7 | 2.9 | 5.4 | 6.4 | 4.1 | 3.6 |
| Black | 10.0 | 6.3 | 2.2 | 2.5 | 6.0 | 3.4 |
| Hispanic | 6.3 | 2.4 | 1.6 | 1.1 | 4.0 | 1.3 |
| Other | 4 . I | 3.4 | 2.3 | 2.6 | 3.5 | 2.2 |
| Grade | | | | | | |
| 6 | 5.7 | 3.3 | 2.5 | 2.6 | 4.0 | 2.2 |
| 7 | 7.5 | 2.7 | 1.3 | 1.4 | 4.5 | 1.7 |
| 8 | 4.8 | 3.4 | 2.3 | 2.7 | 3.5 | 1.8 |
| Total | 6.2 | 1.9 | 2.4 | 1.5 | 4.3 | 1.2 |
| lad ≥ 3 sex partners | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 9.0 | 5.4 | 6.3 | 6.4 | 7.6 | 4.1 |
| Black | 9.4 | 5.1 | 1.4 | 1.8 | 5.2 | 2.6 |
| Hispanic | 7.5 | 3.4 | 2.2 | 1.5 | 4.9 | 1.9 |
| Other | 3.5 | 2.9 | 4.4 | 4 . l | 3.9 | 2.8 |
| Grade | | | | | | |
| 6 | 4.5 | 4.3 | 2.5 | 2.7 | 3.4 | 2.5 |
| 7 | 7.8 | 3.1 | 1.8 | 1.6 | 4.8 | 1.9 |
| 8 | 11.5 | 4.2 | 3.2 | 2.4 | 7.3 | 2.7 |
| Total | 8.0 | 2.5 | 2.9 | 1.6 | 5.4 | 1.6 |

^{*95%} confidence interval.

 $^{^{\}dagger}Non\text{-}Hispanic.$

Table 131. Percentage of Middle School Students Who Had Used a Condom During Last Sexual Intercourse* and Who Had Ever Been Taught About Acquired Immunodeficiency Syndrome (AIDS) or Human Immunodeficiency Virus (HIV) Infection in School, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|-------------------------|---------------------|---------------------|--------|--------|-------|--------|
| Question | <u></u> % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Used condom during last | sexual intercourse | <u> </u> | | | | |
| Race/Ethnicity | | | | | | |
| White§ | 1 | _ | _ | _ | _ | _ |
| Black [§] | _ | _ | _ | _ | | _ |
| Hispanic | 64.3 | 9.7 | _ | _ | 65.0 | 8.5 |
| Other | _ | | _ | _ | | _ |
| Grade | | | | | | |
| 6 | _ | _ | _ | _ | _ | _ |
| 7 | _ | _ | _ | _ | 73.1 | 10.7 |
| 8 | _ | _ | _ | _ | 70.8 | 10.3 |
| Total | 67.3 | 8.9 | _ | _ | 67.5 | 7.7 |
| Had been taught in scho | ol about AIDS or HI | / infection | | | | |
| Race/Ethnicity | | | | | | |
| White | 62.7 | 12.4 | 76.4 | 7.8 | 69.8 | 8.0 |
| Black | 71.1 | 9.9 | 67.9 | 9.5 | 69.4 | 7.6 |
| Hispanic | 58.0 | 7.3 | 57.9 | 7.9 | 57.8 | 7.0 |
| Other | 59.4 | 11.6 | 53.0 | 15.8 | 56.1 | 10.3 |
| Grade | | | | | | |
| 6 | 37.0 | 8.9 | 39.7 | 11.1 | 38.3 | 9.2 |
| 7 | 75.6 | 5.5 | 74.2 | 6.3 | 74.9 | 4.8 |
| 8 | 74.6 | 7.0 | 74.2 | 6.7 | 74.4 | 5.5 |
| Total | 61.4 | 7.2 | 62.7 | 6.1 | 61.9 | 6.3 |

^{*}Among the 15.4% of students who ever had sexual intercourse.

^{†95%} confidence interval.

[§]Non-Hispanic.

¹Not available.

Table 14I. Percentage of Middle School Students Who Had Participated in Sufficient Vigorous Physical Activity,* Who Had Played on Any Sports Teams,† and Who Had Watched ≥ 3 Hours per Day of TV,§ by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | M | Male | | Female | | Total | |
|--------------------------------|---------------------|---------------------|------|--------|------|--------|--|
| Question | % | CI ¹ (±) | % | CI (±) | % | CI (±) | |
| Participated in sufficient vi | gorous physical act | | | | | | |
| Race/Ethnicity | | | | | | | |
| White** | 70.9 | 10.6 | 71.6 | 10.2 | 71.4 | 7.8 | |
| Black** | 63.9 | 8.9 | 54.4 | 10.0 | 59.0 | 7.0 | |
| Hispanic | 66.1 | 6.8 | 66.0 | 5.4 | 66.0 | 5.0 | |
| Other | 71.5 | 8.6 | 65.9 | 9.8 | 68.0 | 7.1 | |
| Grade | | | | | | | |
| 6 | 58.7 | 9.0 | 55.8 | 9.8 | 57.3 | 6.7 | |
| 7 | 68.3 | 5.0 | 66.3 | 6.3 | 67.3 | 4.6 | |
| 8 | 74.8 | 8.1 | 72.8 | 8.4 | 73.8 | 7.1 | |
| Total | 66.9 | 4.8 | 64.2 | 5.1 | 65.6 | 4.0 | |
| Played on any sports teams | 3 | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 52. I | 9.5 | 44.6 | 9.3 | 48.1 | 6.9 | |
| Black | 58.8 | 9.6 | 41.8 | 8.6 | 49.9 | 6.7 | |
| Hispanic | 47.4 | 5.4 | 36.2 | 5.0 | 42.1 | 3.5 | |
| Other | 48. I | 10.0 | 31.2 | 9.6 | 40.2 | 6.6 | |
| Grade | | | | | | | |
| 6 | 49.3 | 7.6 | 34.9 | 9.2 | 42.4 | 5.4 | |
| 7 | 47.6 | 7.0 | 42.5 | 5.2 | 45.I | 4.6 | |
| 8 | 54.3 | 7.3 | 38.5 | 5.7 | 46.4 | 4.7 | |
| Total | 50.3 | 4.1 | 38.7 | 3.9 | 44.6 | 2.6 | |
| Watched \geq 3 hours per day | y of TV | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 50.8 | 8.9 | 46.3 | 11.2 | 48.9 | 7.2 | |
| Black | 54.5 | 10.0 | 66.1 | 8.3 | 60.4 | 5.6 | |
| Hispanic | 59.3 | 4.3 | 52.4 | 4.2 | 56.1 | 3.2 | |
| Other | 58.4 | 10.3 | 44.5 | 9.9 | 51.1 | 6.4 | |
| Grade | | | | | | | |
| 6 | 54.9 | 5.1 | 48.7 | 6.3 | 52.3 | 4.1 | |
| 7 | 57.9 | 6.4 | 53.1 | 5.1 | 55.5 | 4.1 | |
| 8 | 56.5 | 5.3 | 59.2 | 8.8 | 57.8 | 5.7 | |
| Total | 56.7 | 3.1 | 54.1 | 3.7 | 55.6 | 2.6 | |

^{*}Exercised or participated in physical activities that made them sweat and breathe hard for ≥ 20 minutes on ≥ 3 of the 7 days preceding the survey (e.g., basketball, soccer, running, swimming laps, fast bicycling, fast dancing, or similar aerobic activities).

[†]Run by their school or community groups.

[§]On an average school day.

^{195%} confidence interval.

^{**}Non-Hispanic.

Table 151. Percentage of Middle School Students Who Were Enrolled in Physical Education (PE) Class* and Attended PE Class Daily,† by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|---------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Were enrolled in PE class | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 88.2 | 8.5 | 94.3 | 4.7 | 91.2 | 5.9 |
| Black ¹ | 89.0 | 6.5 | 88.4 | 5.8 | 88.7 | 5.3 |
| Hispanic | 88.9 | 3.3 | 92.1 | 2.1 | 90.4 | 2.1 |
| Other | 85.4 | 8.6 | 90.3 | 11.4 | 87.6 | 8.2 |
| Grade | | | | | | |
| 6 | 82.5 | 7.3 | 84.9 | 5.3 | 83.7 | 5.5 |
| 7 | 93.1 | 3.0 | 94.2 | 2.3 | 93.6 | 2.2 |
| 8 | 91.9 | 3.7 | 95.0 | 2.6 | 93.4 | 2.3 |
| Total | 88.7 | 3.2 | 91.4 | 2.1 | 90.0 | 2.4 |
| Attended PE class daily | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 76.7 | 8.9 | 86.0 | 7.0 | 81.3 | 6.2 |
| Black | 76.3 | 9.6 | 76.3 | 8.0 | 76.3 | 6.2 |
| Hispanic | 76.2 | 4.7 | 81.1 | 4.7 | 78.6 | 3.6 |
| Other | 72.3 | 9.1 | 78.8 | 12.3 | 75.5 | 7.7 |
| Grade | | | | | | |
| 6 | 67.0 | 9.0 | 73.9 | 10.5 | 70.6 | 7.6 |
| 7 | 79.8 | 4.4 | 85.3 | 4.4 | 82.5 | 3.7 |
| 8 | 83.7 | 5.9 | 83.5 | 5.8 | 83.6 | 4.8 |
| Total | 76.4 | 4.1 | 80.7 | 4.1 | 78.5 | 3.3 |

^{*}On one or more days in an average week when they were in school.

 $^{^{\}dagger}5$ days in an average week when they were in school.

^{§95%} confidence interval.

¹Non-Hispanic.

Table 16l. Percentage of Middle School Students Who Were at Risk for Becoming* or Were Overweight,† by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|---------------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| At risk for becoming overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 20.4 | 9.4 | 10.0 | 5.3 | 15.2 | 5.3 |
| Black ¹ | 13.5 | 7.4 | 19.3 | 7.3 | 16.3 | 5.2 |
| Hispanic | 18.0 | 3.9 | 22.0 | 4.5 | 19.9 | 2.9 |
| Other | 25.0 | 9.8 | 18.5 | 8.6 | 21.9 | 6.1 |
| Grade | | | | | | |
| 6 | 17.6 | 6.0 | 18.2 | 8.7 | 17.9 | 4.0 |
| 7 | 22.3 | 4.3 | 18.8 | 4.4 | 20.5 | 3.1 |
| 8 | 14.2 | 5.1 | 20.2 | 4.5 | 17.0 | 3.1 |
| Total | 17.9 | 3.1 | 18.9 | 3.2 | 18.4 | 1.9 |
| Overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 22.0 | 8.0 | 12.5 | 8.0 | 17.2 | 6.1 |
| Black | 12.3 | 5.5 | 22.1 | 7.9 | 17.0 | 4.6 |
| Hispanic | 27.7 | 5.4 | 16.6 | 5.7 | 22.4 | 3.9 |
| Other | 16.9 | 9.4 | 17.4 | 6.5 | 17.1 | 6.1 |
| Grade | | | | | | |
| 6 | 24.4 | 6.2 | 22.2 | 14.8 | 23.4 | 8.1 |
| 7 | 25.3 | 5.7 | 14.6 | 3.1 | 19.9 | 3.3 |
| 8 | 18.8 | 7.1 | 15.2 | 4.7 | 17.1 | 4.1 |
| Total | 23.0 | 3.9 | 16.9 | 4.5 | 20.0 | 3.0 |

^{*}Students who were \geq 85th percentile but < 95th percentile for body mass index, by age and sex, based on reference data.

 $^{^{\}dagger}$ Students who were ≥ 95th percentile for body mass index, by age and sex, based on reference data.

^{§95%} confidence interval.

¹Non-Hispanic.

Table 17I. Percentage of Middle School Students Who Described Themselves as Slightly or Very Overweight and Who Were Trying To Lose Weight, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | N | 1ale | Female | | Total | |
|------------------------------------|------|---------|--------|--------------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Described themselves as overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 29.6 | 7.5 | 34.1 | 9.4 | 32.2 | 5.7 |
| Black [†] | 10.1 | 5.4 | 28.8 | 7.1 | 19.6 | 5.2 |
| Hispanic | 27.2 | 4.0 | 27.9 | 5.3 | 27.6 | 3.5 |
| Other | 21.2 | 8.2 | 26.8 | 9.8 | 24.8 | 5.3 |
| Grade | | | | | | |
| 6 | 18.3 | 4.2 | 21.7 | 9.0 | 20.4 | 4.4 |
| 7 | 25.7 | 4.8 | 31.9 | 4.9 | 28.7 | 3.4 |
| 8 | 28.6 | 6.6 | 35.6 | 4.9 | 32.0 | 4.5 |
| Total | 24.3 | 3.1 | 29.2 | 4 . I | 26.9 | 2.7 |
| Were trying to lose weight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 44.8 | 9.2 | 50.7 | 10.1 | 48.0 | 7.6 |
| Black | 28.9 | 9.6 | 49.3 | 9.0 | 39.3 | 6.9 |
| Hispanic | 43.4 | 4.8 | 55.8 | 5.2 | 49.4 | 3.2 |
| Other | 34.0 | 8.5 | 43.6 | 13.1 | 38.9 | 7.8 |
| Grade | | | | | | |
| 6 | 38.6 | 5.7 | 47.5 | 8.1 | 43.3 | 3.4 |
| 7 | 41.8 | 4.8 | 53.9 | 4.6 | 47.7 | 3.9 |
| 8 | 41.7 | 5.2 | 59.1 | 6.2 | 50.2 | 3.5 |
| Total | 40.5 | 3.1 | 52.9 | 3.9 | 46.7 | 2.3 |

^{*95%} confidence interval.

[†]Non-Hispanic.

Table 181. Percentage of Middle School Students Who Had Ever Engaged in Healthy Behaviors Associated with Weight Control To Lose Weight or To Keep from Gaining Weight, by Sex, Race/Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| Question | | Male | | Female | | Total | |
|----------------------------|--------------------------|-------------------|----------------|-----------------|-------|--------|--|
| | % | CI*(±) | % | CI (±) | % | CI (±) | |
| Ate less food, fewer calor | ries, or foods low in fa | at to lose weight | or to keep fro | om gaining weig | ht | | |
| Race/Ethnicity | | | | | | | |
| White [†] | 35.7 | 8.4 | 54.0 | 9.2 | 44.9 | 6.0 | |
| Black [†] | 41.0 | 8.8 | 53.9 | 9.7 | 47.6 | 7.2 | |
| Hispanic | 40.3 | 4.1 | 52.7 | 5.5 | 46.4 | 3.7 | |
| Other | 43.4 | 8.7 | 36.6 | 11.1 | 40.7 | 6.5 | |
| Grade | | | | | | | |
| 6 | 39.9 | 7.8 | 50.4 | 10.1 | 45.3 | 6.7 | |
| 7 | 41.1 | 4.2 | 51.9 | 4.7 | 46.4 | 3.0 | |
| 8 | 37.1 | 5.3 | 56.8 | 5.3 | 46.8 | 3.5 | |
| Total | 39.5 | 3.5 | 52.6 | 4.2 | 46. I | 2.8 | |
| Exercised to lose weight | or to keep from gain | ing weight | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 56.6 | 11.5 | 70.7 | 8.7 | 63.7 | 6.8 | |
| Black | 56.9 | 9.2 | 61.0 | 9.2 | 59.0 | 6.5 | |
| Hispanic | 64.4 | 4.8 | 72 . I | 5.2 | 68. I | 3.9 | |
| Other | 58.9 | 10.0 | 62.5 | 9.4 | 61.0 | 7.1 | |
| Grade | | | | | | | |
| 6 | 55.5 | 7.0 | 64.1 | 8.6 | 60.0 | 7.0 | |
| 7 | 64.0 | 5.3 | 69.9 | 5.1 | 66.9 | 3.0 | |
| 8 | 66.0 | 5.8 | 76.0 | 5.4 | 71.0 | 3.5 | |
| Total | 61.4 | 3.7 | 69. I | 3.9 | 65.3 | 3.0 | |

^{*95%} confidence interval.

[†]Non-Hispanic.

Table 191. Percentage of Middle School Students Who Had Ever Engaged in Unhealthy Behaviors Associated with Weight Control To Lose Weight or To Keep from Gaining Weight, by Sex, Race/ Ethnicity, and Grade—San Bernardino USD, CA, Middle School Youth Risk Behavior Survey, 2003

| | | Male | | Female | | Total | |
|---------------------------|------------------------|-------------------|--------------|--------------------|------|--------|--|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) | |
| Went without eating for | ≥ 24 hours to lose w | eight or to keep | from gaining | | | | |
| Race/Ethnicity | | | | | | | |
| White [†] | 12.4 | 6.2 | 24.7 | 8.6 | 18.4 | 5.3 | |
| Black [†] | 17.9 | 6.9 | 22.2 | 6. l | 20.1 | 4.1 | |
| Hispanic | 14.1 | 3.6 | 20.2 | 4.7 | 17.1 | 2.9 | |
| Other | 19.3 | 8.5 | 12.6 | 5.6 | 16.2 | 4.9 | |
| Grade | | | | | | | |
| 6 | 12.6 | 6.3 | 15.8 | 6.5 | 14.3 | 4.3 | |
| 7 | 17.8 | 4.3 | 20.7 | 4.0 | 19.2 | 3.5 | |
| 8 | 13.5 | 3.8 | 27.5 | 5.7 | 20.4 | 3.5 | |
| Total | 14.6 | 3.1 | 21.0 | 3.2 | 17.9 | 2.3 | |
| Took diet pills, powders, | or liquids to lose wei | ght or to keep fr | om gaining w | eight [§] | | | |
| Race/Ethnicity | | | | | | | |
| White | 3.0 | 2.6 | 9.0 | 5.4 | 5.9 | 2.8 | |
| Black | 8.7 | 5.5 | 5.3 | 4 . I | 6.9 | 3.8 | |
| Hispanic | 6.1 | 2.2 | 8.3 | 2.7 | 7.2 | 1.8 | |
| Other | 6.6 | 5.6 | 7.0 | 5.0 | 7.0 | 3.9 | |
| Grade | | | | | | | |
| 6 | 5.3 | 4.4 | 2.9 | 2.2 | 4.1 | 2.3 | |
| 7 | 5.7 | 2.2 | 7.0 | 2.5 | 6.3 | 1.8 | |
| 8 | 7.4 | 3.2 | 13.7 | 4.9 | 10.5 | 3.2 | |
| Total | 6.0 | 2.0 | 7.7 | 2.2 | 6.8 | 1.6 | |
| omited or took laxative | s to lose weight or to | keep from gain | ing weight | | | | |
| Race/Ethnicity | | | | | | | |
| White | 1.9 | 2.3 | 6.8 | 4.2 | 4.3 | 2.3 | |
| Black | 8.5 | 6.0 | 6.0 | 4.3 | 7.2 | 3.6 | |
| Hispanic | 5.0 | 2.1 | 6.5 | 2.5 | 5.9 | 1.9 | |
| Other | 7.1 | 5.5 | 10.2 | 7.9 | 8.8 | 4.5 | |
| Grade | | | | | | | |
| 6 | 5.0 | 3.6 | 3.3 | 3.0 | 4.4 | 1.9 | |
| 7 | 5.1 | 2.8 | 6.3 | 2.5 | 5.7 | 1.9 | |
| 8 | 5.7 | 3.0 | 10.2 | 4.2 | 7.9 | 2.9 | |
| Total | 5.2 | 1.9 | 6.5 | 2.0 | 5.9 | 1.4 | |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Without a doctor's advice.